Journal of Mathematical Research with Applications Sept., 2014, Vol. 34, No. 5, pp. 505–516 DOI:10.3770/j.issn:2095-2651.2014.05.001 Http://jmre.dlut.edu.cn

# Laplacian Spectral Characterization of a Kind of Unicyclic Graphs

Luhua WANG, Ligong WANG\*

Department of Applied Mathematics, School of Science, Northwestern Polytechnical University, Shaanxi 710072, P. R. China

**Abstract** Let  $H(n; q, n_1, n_2, n_3, n_4)$  be a unicyclic graph with n vertices containing a cycle  $C_q$  and four hanging paths  $P_{n_1+1}$ ,  $P_{n_2+1}$ ,  $P_{n_3+1}$  and  $P_{n_4+1}$  attached at the same vertex of the cycle. In this paper, it is proved that all unicyclic graphs  $H(n; q, n_1, n_2, n_3, n_4)$  are determined by their Laplacian spectra.

Keywords Laplacian spectrum; unicyclic graphs; Laplacian matrix.

MR(2010) Subject Classification 05C50; 15A18

#### 1. Introduction

In this paper, we assume that all graphs are simple and undirected. Let G = (V(G), E(G))be a graph with vertex set  $V(G) = \{v_1, v_2, \ldots, v_n\}$  and edge set E(G). Let the matrix A(G) be the (0, 1)-adjacency matrix of G and  $d_i = d(v_i) = d_i(G)$  the degree of the vertex  $v_i$ . Assume that  $d_1 \ge d_2 \ge \cdots \ge d_n$ . The matrix L(G) = D(G) - A(G) is called the Laplacian matrix of G, where  $D(G) = \text{diag}(d_1, d_2, \ldots, d_n)$  denotes the diagonal matrix of vertex degrees of G. The polynomial  $\phi(G) = \phi(G, x) = \det(xI_n - L(G)) = l_0x^n + l_1x^{n-1} + \cdots + l_n$  is defined as the Laplacian characteristic polynomials of the graph G, where  $I_n$  is the  $n \times n$  identity matrix. Since A(G) and L(G) are real and symmetric, their eigenvalues are real numbers and called the adjacency and the Laplacian eigenvalues of G, respectively. Assume that  $\lambda_1 \ge \lambda_2 \ge \cdots \ge \lambda_n$  and  $\mu_1 \ge \mu_2 \ge \cdots \ge \mu_n (= 0)$  are the adjacency and the Laplacian eigenvalues of G. The multiset of eigenvalues of A(G) (L(G)) is called the adjacency (Laplacian) spectrum of G. Two graphs are said to be adjacency (Laplacian) cospectral if they have equal adjacency (Laplacian) spectrum. A graph is said to be determined by the adjacency (Laplacian) spectrum if there is no other non-isomorphic graph with the same adjacency (Laplacian) spectrum.

Which graphs are determined by their (Laplacian) spectra is an interesting problem in the theory of graph spectra. Characterizing such graphs seems also to be a difficult problem. Recently, this problem has attracted some researchers' attention. Up until now, only some graphs with special structure are proved to be determined by their (Laplacian) spectra. Some results

Received November 14, 2013; Accepted June 19, 2014

Supported by the National Natural Science Foundation of China (Grant No. 11171273) and Graduate Starting Seed Fund of Northwestern Polytechnical University (Grant No. Z2014173).

\* Corresponding author

E-mail address: luhuawang126@163.com (Luhua WANG); lgwangmath@163.com (Ligong WANG)

on these special graphs determined by their adjacency or Laplacian spectra can be found in [10, 15, 16, 18, 24, 26, 27] or [1, 10, 17-20, 22, 23, 25, 28], respectively. The reader can also consult the books [3, 5] and the surveys [6, 7].

Let  $P_n$  and  $C_n$  be the path and cycle with n vertices, respectively. Let a kind of unicyclic graphs  $H(n; q, n_1, n_2, ..., n_t)$  with n vertices be shown in Figure 1. They are some unicyclic graphs containing a cycle  $C_q$  and t hanging paths  $P_{n_1+1}, P_{n_2+1}, P_{n_3+1}, ..., P_{n_t+1}$  attached at the same vertex of the cycle. For the case t = 1, Haemers et al. in [10] have proved all lollipop graphs H(n; q, n - q) to be determined by their Laplacian spectra. For the case t = 2, Liu et al. in [18] has proved all graphs  $H(n; q, n_1, n_2)$  to be determined by their Laplacian spectra. For the case t = 3, Lu et al. in [20] have proved all graphs  $H(n; q, n_1, n_2, n_3)$  to be determined by their Laplacian spectra. In this paper, we prove all unicyclic graphs  $H(n; q, n_1, n_2, n_3, n_4)$  (shown in Figure 2) to be determined by their Laplacian spectra.



Figure 1 Graph  $H(n; q, n_1, n_2, \ldots, n_t)$ 



Figure 2 Graph  $H(n;q,n_1,n_2,n_3,n_4)$ 

#### 2. Preliminaries

In this section, some known lemmas about the (Laplacian) spectrum of a graph are given, and these lemmas will play important roles throughout the paper.

**Lemma 2.1** ([6,21]) For the Laplacian matrix of a graph G, the following results can be deduced from the Laplacian spectrum.

- (i) The number of vertices.
- (ii) The number of edges.
- (iii) The number of components.
- (iv) The number of spanning trees.
- (v) The sum of the squares of degrees of vertices.

**Lemma 2.2** ([5]) Assume that N is an  $n \times n$  symmetric matrix, its eigenvalues are  $\alpha_1 \ge \alpha_2 \ge \cdots \ge \alpha_n$ . The eigenvalues of the principal submatrix of order m of N are  $\alpha'_1 \ge \alpha'_2 \ge \cdots \ge \alpha'_m$ , then  $\alpha_i \ge \alpha'_i \ge \alpha_{n-m+i}$ , i = 1, 2, ..., m.

**Lemma 2.3** ([4]) Let  $A = [a_{ij}]$  be a matrix of order n. Let

$$r_i = \sum_{j=1}^n |a_{ij}|, \quad 1 \le i \le n$$

be the sum of the absolute values of the entries in row i of A. Then

$$\rho(A) \le \max(r_1, r_2, \dots, r_n)$$

where  $\rho(A)$  is the maximum of the absolute values of all eigenvalues of A. A similar inequality holds for the sum of the absolute values of the entries in columns of A.

**Lemma 2.4** ([21]) Let G be a graph with n vertices and m edges and let  $\deg(G) = (d_1, d_2, \dots, d_n)$  be its non-increasing degree sequence. Then the first four coefficients in  $\phi(G, x)$  are

$$l_0 = 1, l_1 = -2m, l_2 = 2m^2 - m - \frac{1}{2} \sum_{i=1}^n d_i^2,$$
  
$$l_3 = \frac{1}{3} (-4m^3 + 6m^2 + 3m \sum_{i=1}^n d_i^2 - \sum_{i=1}^n d_i^3 - 3\sum_{i=1}^n d_i^2 + 6n_G(C_3)),$$

where  $n_G(C_3)$  is the number of triangles in G.

**Lemma 2.5** ([12, 14]) Let G be a graph with  $V(G) \neq \emptyset$  and  $E(G) \neq \emptyset$ . Then

$$d_1 + 1 \le \mu_1 \le \max\{\frac{d_i(d_i + m_i) + d_j(d_j + m_j)}{d_i + d_j}, v_i v_j \in E(G)\}$$

where  $d_1$  denotes the maximum vertex degree of G and  $m_i$  denotes the average of the degrees of the vertices adjacent to vertex  $v_i$  in G.

**Lemma 2.6** ([13]) Let G be a connected graph with  $n \ge 3$  vertices, and let  $\deg(G) = (d_1, d_2, \ldots, d_n)$  be its non-increasing degree sequence. Then  $\mu_2 \ge d_2$ .

**Lemma 2.7** ([2]) The coefficients  $l_i$  of the polynomial  $\phi(G, x)$  are given by the formula

$$(-1)^i l_i = \sum p(\Phi), \quad 1 \le i \le n,$$

where the right hand of the equation is the summation of all sub-forests  $\Phi$  of G which have i edges, and  $p(\Phi)$  denotes the product of the numbers of vertices in the components of the forest  $\Phi$ .

**Lemma 2.8** ([8]) Let  $B_n$  be the matrix of order n obtained from  $L(P_{n+1})$  by deleting the row and column corresponding to some end vertices of  $P_{n+1}$ , and let  $H_n$  be the matrix of order nobtained from  $L(P_{n+2})$  by deleting the rows and columns corresponding to the two end vertices of  $P_{n+2}$ . Set  $\phi(P_0) = 0, \phi(B_0) = 1, \phi(H_0) = 1$ . Then we have the following conclusions.

- (i)  $\phi(P_{n+1}) = (x-2)\phi(P_n) \phi(P_{n-1}), (n \ge 1).$
- (ii)  $x\phi(B_n) = \phi(P_{n+1}) + \phi(P_n).$
- (iii)  $\phi(P_n) = x(\phi(H_{n-1})) \ (n \ge 1).$
- (iv)  $\phi(C_n) = \frac{1}{x}\phi(P_{n+1}) \frac{1}{x}\phi(P_{n-1}) + 2(-1)^{n+1} \ (n \ge 3, x \ne 0).$
- From Lemma 2.8, if x = 4, then we can get the following lemma.

**Lemma 2.9** ([20]) (i)  $\phi(P_n; 4) = 4n$ .

(ii) 
$$\phi(B_n; 4) = 2n + 1.$$

- (iii)  $\phi(H_n; 4) = n + 1.$
- (iv)  $\phi(C_n; 4) = 2 + 2(-1)^{n+1}$ .

From Lemma 2.8, if  $x \neq 4$ , then we can get the following lemma.

**Lemma 2.10** For  $\phi(G) = \phi(G, x)$  of a graph G, if  $x \neq 4$ , and y satisfies the equation  $y^2 - (x - 2)y + 1 = 0$ . Then we have the following results.

(i)  $\phi(P_n) = \frac{(y+1)(y^{2n}-1)}{y^{n+1}-y^n}$ . (ii)  $\phi(B_n) = \frac{y^{2n+1}-1}{y^{n+1}-y^n}$ . (iii)  $\phi(H_n) = \frac{y^{2n+2}-1}{y^{n+2}-y^n}$ . (iv)  $\phi(C_n) = y^n + \frac{1}{y^n} + 2(-1)^{n+1}$ .

If v is a vertex of G, let  $L_v(G)$  be the principal submatrix of L(G) obtained by deleting the row and column corresponding to vertex v from L(G).

**Lemma 2.11**([9]) Let  $G_1$  and  $G_2$  be two vertex-disjoint graphs and  $G = G_1 u : vG_2$  be a graph obtained by joining the vertex u of the graph  $G_1$  to the vertex v of the graph  $G_2$  by an edge. Then  $\phi(G) = \phi(G_1)\phi(G_2) - \phi(G_1)\phi(L_v(G_2)) - \phi(G_2)\phi(L_u(G_1))$ .

**Lemma 2.12** ([18]) Let G be a connected unicyclic graph with n vertices and its cycle  $C_q$ . If G' is Laplacian cospectral to G, then G' must be a connected unicyclic graph with n vertices and one cycle  $C_q$ . Moreover,

$$\sum_{i=1}^{n} d_i(G)^3 = \sum_{i=1}^{n} d_i(G')^3.$$

## **3.** Laplacian spectral characterization of graphs $H(n; q, n_1, n_2, n_3, n_4)$

In this section, we prove that no two non-isomorphic graphs of the form  $H(n; q, n_1, n_2, n_3, n_4)$  are Laplacian cospectral, and the graph  $H(n; q, n_1, n_2, n_3, n_4)$  is determined by its Laplacian spectrum. To prove Theorems 3.3 and 3.4, we first prove the following lemmas.

**Lemma 3.1** The antepenultimate coefficient  $l_{n-2}$  of  $\phi(H(n;q,n_1,n_2,n_3,n_4),x)$  is given by

$$(-1)^{n-2}l_{n-2} = q[\frac{1}{6}(n_1 + n_2 + n_3 + n_4)^3 - (n_1n_2n_3 + n_1n_2n_4 + n_1n_3n_4 + n_2n_3n_4) + \frac{1}{2}q(n_1 + n_2 + n_3 + n_4)^2 + (1 - q)(n_1n_2 + n_1n_3 + n_1n_4 + n_2n_3 + n_2n_4 + n_3n_4) + (\frac{1}{2}q - \frac{1}{6})(n_1 + n_2 + n_3 + n_4)] + \sum p(\Phi),$$

where the summation is over all sub-forests  $\Phi$  of  $H(n;q,n_1,n_2,n_3,n_4)$  which have n-2 edges obtained by deleting two edges both from  $C_q$  in  $H(n;q,n_1,n_2,n_3,n_4)$ .

**Proof** Let  $G = H(n; q, n_1, n_2, n_3, n_4)$ . Since G has n edges, the (n - 2)-subforests of G can be obtained from G by deleting two edges in which at least one comes from the cycle  $C_q$ . Then by Lemma 2.7, we can get that

$$(-1)^{n-2}l_{n-2} = q \sum_{i=0}^{n_1-1} (q+n_2+n_3+n_4+i)(n_1-i) + q \sum_{i=0}^{n_2-1} (q+n_1+n_3+n_4+i)(n_2-i) + q \sum_{i=0}^{n_3-1} (q+n_1+n_2+n_4+i)(n_3-i) + q \sum_{i=0}^{n_4-1} (q+n_1+n_2+n_3+i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_1+n_2+n_3+i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_1+n_2+n_3+i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_1+n_2+n_4+i)(n_3-i) + q \sum_{i=0}^{n_4-1} (q+n_1+n_2+n_4+i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_4+i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_4+i)(n_4-i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_4+i)(n_4-i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_4+i)(n_4-i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_4+i)(n_4-i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_4+i)(n_4-i)(n_4-i)(n_4-i) + q \sum_{i=0}^{n_4-1} (q+n_4-i)(n_4$$

 $\label{eq:laplacian} Laplacian\ spectral\ characterization\ of\ a\ kind\ of\ unicyclic\ graphs$ 

$$\sum p(\Phi),\tag{1}$$

where the summation is over all sub-forests  $\Phi$  of  $G = H(n; q, n_1, n_2, n_3, n_4)$  which have n - 2 edges obtained by deleting two edges both from  $C_q$  in G.

Because

$$\sum_{i=0}^{n_1-1} (q+n_2+n_3+n_4+i)(n_1-i)$$

$$=\sum_{i=0}^{n_1-1} (q+n_2+n_3+n_4)n_1 + \sum_{i=0}^{n_1-1} (n_1-q-n_2-n_3-n_4)i - \sum_{i=0}^{n_1-1} i^2$$

$$= (q+n_2+n_3+n_4)n_1^2 + (n_1-q-n_2-n_3-n_4)\frac{n_1(n_1-1)}{2} - \frac{1}{6}n_1(n_1-1)(2n_1-1)$$

$$= \frac{1}{6}n_1^3 + \frac{1}{2}n_1^2n_2 + \frac{1}{2}n_1^2n_3 + \frac{1}{2}n_1^2n_4 + \frac{1}{2}n_1n_2 + \frac{1}{2}n_1n_3 + \frac{1}{2}n_1n_4 + \frac{1}{2}n_1q - \frac{1}{6}n_1 + \frac{1}{2}n_1^2q,$$
(2)

similarly, we obtain the following results:

$$\sum_{i=0}^{n_2-1} (q+n_1+n_3+n_4+i)(n_2-i)$$
  
=  $\frac{1}{6}n_2^3 + \frac{1}{2}n_2^2n_1 + \frac{1}{2}n_2^2n_3 + \frac{1}{2}n_2^2n_4 + \frac{1}{2}n_2n_1 + \frac{1}{2}n_2n_3 + \frac{1}{2}n_2n_4 + \frac{1}{2}n_2q - \frac{1}{6}n_2 + \frac{1}{2}n_2^2q,$  (3)

$$\sum_{i=0}^{n_3-1} (q+n_1+n_2+n_4+i)(n_3-i) \\ = \frac{1}{6}n_3^3 + \frac{1}{2}n_3^2n_1 + \frac{1}{2}n_3^2n_2 + \frac{1}{2}n_3^2n_4 + \frac{1}{2}n_3n_1 + \frac{1}{2}n_3n_2 + \frac{1}{2}n_3n_4 + \frac{1}{2}n_3q - \frac{1}{6}n_3 + \frac{1}{2}n_3^2q,$$
(4)

$$\sum_{i=0}^{n_4-1} (q+n_1+n_2+n_3+i)(n_4-i)$$

$$= \frac{1}{6}n_4^3 + \frac{1}{2}n_4^2n_1 + \frac{1}{2}n_4^2n_2 + \frac{1}{2}n_4^2n_3 + \frac{1}{2}n_4n_1 + \frac{1}{2}n_4n_2 + \frac{1}{2}n_4n_3 + \frac{1}{2}n_4q - \frac{1}{6}n_4 + \frac{1}{2}n_4^2q.$$
(5)

Then, substituting (2)–(5) into (1), we have  $\frac{1}{1}$ 

$$(-1)^{n-2}l_{n-2} = q[\frac{1}{6}(n_1 + n_2 + n_3 + n_4)^3 - (n_1n_2n_3 + n_1n_2n_4 + n_1n_3n_4 + n_2n_3n_4) + \frac{1}{2}q(n_1 + n_2 + n_3 + n_4)^2 + (1 - q)(n_1n_2 + n_1n_3 + n_1n_4 + n_2n_3 + n_2n_4 + n_3n_4) + (\frac{1}{2}q - \frac{1}{6})(n_1 + n_2 + n_3 + n_4)] + \sum p(\Phi),$$

where the summation is over all sub-forests  $\Phi$  of  $G = H(n; q, n_1, n_2, n_3, n_4)$  which have n - 2 edges obtained by deleting two edges both from  $C_q$  in G.  $\Box$ 

**Lemma 3.2** Let G be the graph  $H(n; q, n_1, n_2, n_3, n_4)$  as shown in Figure 2. Then we have the following result:

$$\phi(G,4) = (32 - 128q - 32(-1)^q)n_1n_2n_3n_4 + (16 - 48q - 16(-1)^q) \cdot (n_1n_2n_3 + n_1n_2n_4 + n_1n_3n_4 + n_2n_3n_4) + (8 - 16q - 8(-1)^q) \cdot (n_1n_2 + n_1n_3 + n_1n_4 + n_2n_3 + n_2n_4 + n_3n_4) + (4 - 4q - 4(-1)^q) \cdot (n_1 + n_2 + n_3 + n_4) + 2 - 2(-1)^q.$$

**Proof** Firstly, we assume that  $G_2 = H(n - n_1; q, n_2, n_3, n_4)$ ,  $G_3 = H(n - n_1 - n_2; q, n_3, n_4)$  and  $G_4 = H(n - n_1 - n_2 - n_3; q, n_4)$ . Then, from Lemma 2.11, we obtain

$$\begin{split} \phi(G,x) &= \phi(G_2,x)\phi(P_{n_1},x) - \phi(G_2,x)\phi(B_{n_1-1},x) - \\ &\phi(P_{n_1},x)\phi(H_{q-1},x)\phi(B_{n_2},x)\phi(B_{n_3},x)\phi(B_{n_4},x). \end{split}$$
(6)  
$$\phi(G_2,x) &= \phi(G_3,x)\phi(P_{n_2},x) - \phi(G_3,x)\phi(B_{n_2-1},x) - \\ &\phi(P_{n_2},x)\phi(H_{q-1},x)\phi(B_{n_3},x)\phi(B_{n_4},x). \\ \phi(G_3,x) &= \phi(G_4,x)\phi(P_{n_3},x) - \phi(G_4,x)\phi(B_{n_3-1},x) - \\ &\phi(P_{n_3},x)\phi(H_{q-1},x)\phi(P_{n_4},x). \\ \phi(G_4,x) &= \phi(C_q,x)\phi(P_{n_4},x) - \phi(C_q,x)\phi(B_{n_4-1},x) - \\ &\phi(H_{q-1},x)\phi(B_{n_4},x). \end{split}$$

When x = 4, by Lemma 2.9, we obtain the following results:

$$\phi(G_4, 4) = 4n_4 - 4(-1)^q n_4 - 4qn_4 + 2 - 2(-1)^q, \tag{7}$$

$$\phi(G_{3},4) = 8n_{3}n_{4} - 8(-1)^{q}n_{3}n_{4} - 16n_{3}n_{4}q + 4n_{3} + 4n_{4} - 4(-1)^{q}n_{3} - 4(-1)^{q}n_{4} - 4n_{4}q - 4n_{3}q + 2 - 2(-1)^{q},$$

$$\phi(G_{2},4) = 16n_{2}n_{3}n_{4} - 16(-1)^{q}n_{2}n_{3}n_{4} - 48n_{2}n_{3}n_{4}q + 8n_{2}n_{3} + 8n_{2}n_{4} + 8n_{3}n_{4} - 8(-1)^{q}n_{2}n_{3} - 8(-1)^{q}n_{2}n_{4} - 8(-1)^{q}n_{2}n_{4} - 8(-1)^{q}n_{3}n_{4} - 16n_{2}n_{4}q - 16n_{2}n_{3}q - 16n_{3}n_{4}q + 4n_{2} + 4n_{3} + 4n_{4} - 4(-1)^{q}n_{2} - 4(-1)^{q}n_{3} - 4(-1)^{q}n_{4} - 8(-1)^{q}n_{4} - 8(-1$$

$$4n_2q - 4n_3q - 4n_4q + 2 - 2(-1)^q. (9)$$

Then, substituting (7)-(9) into (6), we can get that

$$\phi(G,4) = (32 - 128q - 32(-1)^q)n_1n_2n_3n_4 + (16 - 48q - 16(-1)^q) \cdot (n_1n_2n_3 + n_1n_2n_4 + n_1n_3n_4 + n_2n_3n_4) + (8 - 16q - 8(-1)^q) \cdot (n_1n_2 + n_1n_3 + n_1n_4 + n_2n_3 + n_2n_4 + n_3n_4) + (4 - 4q - 4(-1)^q) \cdot (n_1 + n_2 + n_3 + n_4) + 2 - 2(-1)^q. \quad \Box$$

$$(10)$$

1

Note that when  $x \neq 4$  and y satisfies the equation  $y^2 - (x - 2)y + 1 = 0$ , by Lemma 2.10, we can get the following expressions of  $\phi(G) = \phi(H(n; q, n_1, n_2, n_3, n_4))$  by using Mathematica:

$$\begin{split} \phi(G_4) &= \frac{1}{(y-1)^2} [-2(-1)^{1+q} y^{1-n_4} + 2(-1)^{1+q} y^{-n_4} - 2(-1)^{1+q} y^{1+n_4} + 2(-1)^{1+q} y^{2+n_4} + y^{-n_4-q} - 2y^{1-n_4-q} + y^{2+n_4-q} + y^{-n_4+q} - 2y^{1+n_4+q} + y^{2+n_4+q}], \end{split} \tag{11}$$

Then, substituting (11)-(13) into (6), we can get that

$$(y-1)^{5}\phi(G) + 1 - 5y + y^{2q} + 3y^{1+2q} + 5y^{4+2n} - y^{5+2n} - 3y^{4+2(n-q)} - y^{5+2(n-q)} + 2(-1)^{1+q}y^{q} - 2(-1)^{1+q}y^{1+q} + 2(-1)^{1+q}y^{4+2n-q} - 2(-1)^{1+q}y^{5+2n-q} = f(n_1, n_2, n_3, n_4; y)$$

$$(14)$$

where  $n = q + n_1 + n_2 + n_3 + n_4$ ,

 $f(n_1, n_2, n_3, n_4; y)$ 

$$\begin{split} &= -4y^{2+2n_1} - 4y^{2+2n_2} - 4y^{2+2n_3} - 4y^{2+2n_4} + y^{2+2n_1+2n_2} + y^{2+2n_1+2n_3} + \\ &y^{2+2n_1+2n_4} + y^{2+2n_2+2n_3} + y^{2+2n_2+2n_4} + y^{2+2n_3+2n_4} + 3y^{3+2n_1+2n_2} + \\ &3y^{3+2n_1+2n_3} + 3y^{3+2n_1+2n_4} + 3y^{3+2n_2+2n_3} + 3y^{3+2n_2+2n_4} + 3y^{3+2n_3+2n_4} - \\ &2y^{3+2n_1+2n_2+2n_3} - 2y^{3+2n_1+2n_2+2n_4} - 2y^{3+2n_1+2n_3+2n_4} - 2y^{3+2n_2+2n_3+2n_4} - \\ &2y^{4+2n_1+2n_2+2n_3} - 2y^{4+2n_1+2n_2+2n_4} - 2y^{4+2n_1+2n_3+2n_4} - 2y^{4+2n_2+2n_3+2n_4} + \\ &2(-1)^{1+q}y^{1+2n_1+q} + 2(-1)^{1+q}y^{1+2n_2+q} + 2(-1)^{1+q}y^{1+2n_3+q} + \\ &2(-1)^{1+q}y^{1+2n_4+q} - 2(-1)^{1+q}y^{2+2n_1+q} - 2(-1)^{1+q}y^{2+2n_2+q} - \\ &2(-1)^{1+q}y^{2+2n_3+q} - 2(-1)^{1+q}y^{2+2n_1+q} - 2(-1)^{1+q}y^{2+2n_2+2n_3+q} - \\ &2(-1)^{1+q}y^{2+2n_2+2n_4+q} - 2(-1)^{1+q}y^{2+2n_3+2n_4+q} + 2(-1)^{1+q}y^{3+2n_1+2n_2+q} + \\ &2(-1)^{1+q}y^{3+2n_1+2n_3+q} + 2(-1)^{1+q}y^{3+2n_1+2n_4+q} + 2(-1)^{1+q}y^{3+2n_1+2n_2+q} + \\ &2(-1)^{1+q}y^{3+2n_1+2n_3+q} + 2(-1)^{1+q}y^{3+2n_3+2n_4+q} - 2(-1)^{1+q}y^{4+2n_1+2n_2+2n_3+q} + \\ &2(-1)^{1+q}y^{4+2n_1+2n_2+2n_4+q} + 2(-1)^{1+q}y^{3+2n_1+2n_3+2n_4+q} - \\ &2(-1)^{1+q}y^{4+2n_1+2n_2+2n_4+q} + 2(-1)^{1+q}y^{3+2n_1+2n_3+2n_4+q} - \\ &2(-1)^{1+q}y^{4+2n_1+2n_2+2n_4+q} + 2(-1)^{1+q}y^{3+2n_1+2n_3+2n_4+q} + \\ &2(-1)^{1+q}y^{3+2n_2+2n_3+2n_4+q} + 2(-1)^{1+q}y^{3+2n_1+2n_3+2n_4+q} + \\ &2(-1)^{1+q}y^{3+2n_2+2n_3+2n_4+q} + 2(-1)^{1+q}y^{3+2n_1+2n_3+2n_4+q} + \\ &2(-1)^{1+q}y^{3+2n_2+2n_3+2n_4+q} + 2y^{1+2n_1+2q} + 2y^{1+2n_3+2q} + \\ &2y^{1+2n_4+2q} + 2y^{2+2n_1+2q} + 2y^{2+2n_3+2q} - 3y^{2+2n_1+2n_3+2q} + \\ &3y^{2+2n_1+2n_2+2q} - 3y^{2+2n_1+2n_3+2q} - 3y^{2+2n_1+2n_4+2q} - 3y^{2+2n_2+2n_3+2q} - \\ &3y^{2+2n_1+2n_4+2q} - y^{3+2n_2+2n_3+2q} - y^{3+2n_1+2n_4+2q} - y^{3+2n_3+2n_4+2q} + \\ &4y^{3+2n_1+2n_2+2n_3+2q} + 4y^{3+2n_1+2n_2+2n_4+2q} + 4y^{3+2n_1+2n_3+2n_4+2q} + \\ &4y^{3+2n_1+2n_2+2n_3+2q} + 4y^{3+2n_1+2n_2+2n_4+2q} + 4y^{3+2n_1+2n_3+2n_4+2q} + \\ &4y^{3+2n_1+2n_2+2n_3+2q} + 2y^{3+2n_2+2n_3+2q} + 2y^{3+2n_3+2n_4+2q} + \\ &4y^{3+2n_2+2n_3+2n_4+2q} + 2y^{3+2n_3+2n_4+2q} + \\ &4y^{3+2n_2+2n_3$$

**Theorem 3.3** No two non-isomorphic graphs of the form  $H(n; q, n_1, n_2, n_3, n_4)$  are Laplacian cospectral.

**Proof** Suppose that  $G' = H(n; q', n'_1, n'_2, n'_3, n'_4)$  is Laplacian cospectral to  $G = H(n; q, n_1, n_2, n_3, n_4)$ . From Lemma 2.12, we have q = q', then  $n_1 + n_2 + n_3 + n_4 = n'_1 + n'_2 + n'_3 + n'_4$ . In the following, we prove that G and G' are isomorphic.

We consider the coefficient  $l_{n-2}$  of  $\phi(G, x)$  and the coefficient  $l'_{n-2}$  of  $\phi(G', x)$ . From Lemma 3.1, we can get  $l_{n-2}$  of  $\phi(G)$  and  $l'_{n-2}$  of  $\phi(G')$ .

$$(-1)^{n-2}l_{n-2} = q[\frac{1}{6}(n_1 + n_2 + n_3 + n_4)^3 - (n_1n_2n_3 + n_1n_2n_4 + n_1n_3n_4 + n_2n_3n_4) + \frac{1}{2}q(n_1 + n_2 + n_3 + n_4)^2 + (1 - q)(n_1n_2 + n_1n_3 + n_1n_4 + n_2n_3 + n_2n_4 + n_3n_4) + (\frac{1}{2}q - \frac{1}{6})(n_1 + n_2 + n_3 + n_4)] + \sum p(\Phi),$$

Laplacian spectral characterization of a kind of unicyclic graphs

$$\begin{split} (-1)^{n-2}l'_{n-2} =& q'[\frac{1}{6}(n'_1+n'_2+n'_3+n'_4)^3 - (n'_1n'_2n'_3+n'_1n'_2n'_4+n'_1n'_3n'_4+n'_2n'_3n'_4) + \\ & \frac{1}{2}q'(n'_1+n'_2+n'_3+n'_4)^2 + (1-q')(n'_1n'_2+n'_1n'_3+n'_1n'_4+n'_2n'_3+ \\ & n'_2n'_4+n'_3n'_4) + (\frac{1}{2}q'-\frac{1}{6})(n'_1+n'_2+n'_3+n'_4)] + \sum p(\Phi'), \end{split}$$

where the summation is over all sub-forests  $\Phi$  (resp.,  $\Phi'$ ) of G(G') with n-2 (or n'-2) edges obtained by deleting two edges both from  $C_q$  in G(G').

Because  $q = q', n_1 + n_2 + n_3 + n_4 = n'_1 + n'_2 + n'_3 + n'_4$ ,  $\sum p(\Phi) = \sum p(\Phi')$ , and  $(-1)^{n-2}l_{n-2} = (-1)^{n-2}l'_{n-2}$ , we have

$$(n_1n_2n_3 + n_1n_2n_4 + n_1n_3n_4 + n_2n_3n_4) - (1-q)(n_1n_2 + n_1n_3 + n_1n_4 + n_2n_3 + n_2n_4 + n_3n_4) = (n'_1n'_2n'_3 + n'_1n'_2n'_4 + n'_1n'_3n'_4 + n'_2n'_3n'_4) - (1-q)(n'_1n'_2 + n'_1n'_3 + n'_1n'_4 + n'_2n'_3 + n'_2n'_4 + n'_3n'_4).$$
(15)

From Lemma 3.2, we can get  $\phi(G, 4)$  and  $\phi(G', 4)$  as follows

$$\begin{split} \phi(G,4) =& (32-128q-32(-1)^q)n_1n_2n_3n_4 + (16-48q-16(-1)^q) \cdot \\ & (n_1n_2n_3+n_1n_2n_4+n_1n_3n_4+n_2n_3n_4) + (8-16q-8(-1)^q) \cdot \\ & (n_1n_2+n_1n_3+n_1n_4+n_2n_3+n_2n_4+n_3n_4) + (4-4q-4(-1)^q) \cdot (n_1+n_2+n_3+n_4) + 2-2(-1)^q, \\ \phi(G',4) =& (32-128q'-32(-1)^{q'})n_1'n_2'n_3'n_4' + (16-48q'-16(-1)^{q'}) \cdot \\ & (n_1'n_2'n_3'+n_1'n_2'n_4'+n_1'n_3'n_4'+n_2'n_3'n_4') + (8-16q'-8(-1)^{q'}) \cdot \\ & (n_1'n_2'+n_1'n_3'+n_1'n_4'+n_2'n_3'+n_2'n_4'+n_3'n_4') + (4-4q'-4(-1)^{q'}) \cdot (n_1'+n_2'+n_3'+n_4') + 2-2(-1)^{q'}. \end{split}$$

Because  $\phi(G, 4) = \phi(G', 4), q = q'$ , we have

$$(4 - 16q - 4(-1)^{q})n_{1}n_{2}n_{3}n_{4} + (2 - 6q - 2(-1)^{q})(n_{1}n_{2}n_{3} + n_{1}n_{2}n_{4} + n_{1}n_{3}n_{4} + n_{2}n_{3}n_{4}) + (1 - 2q - (-1)^{q})(n_{1}n_{2} + n_{1}n_{3} + n_{1}n_{4} + n_{2}n_{3} + n_{2}n_{4} + n_{3}n_{4}) = (4 - 16q - 4(-1)^{q})n_{1}'n_{2}'n_{3}'n_{4}' + (2 - 6q - 2(-1)^{q}) \cdot (n_{1}'n_{2}'n_{3}' + n_{1}'n_{2}'n_{4}' + n_{1}'n_{3}'n_{4}' + n_{2}'n_{3}'n_{4}') + (1 - 2q - (-1)^{q})(n_{1}'n_{2}' + n_{1}'n_{3}' + n_{1}'n_{3}'n_{4}' + n_{2}'n_{3}'n_{4}').$$
(16)

Because the graphs G and G' have the same Laplacian characteristic polynomials, that is,  $\phi(G) = \phi(G')$ , by (14) we have  $f(n_1, n_2, n_3, n_4; y) = f(n'_1, n'_2, n'_3, n'_4; y)$ . Without loss of generality, assume that  $n_1 \leq n_2 \leq n_3 \leq n_4$  and  $n'_1 \leq n'_2 \leq n'_3 \leq n'_4$ , then the smallest degrees of all items of the polynomials  $f(n_1, n_2, n_3, n_4; y)$  and  $f(n'_1, n'_2, n'_3, n'_4; y)$  on variable y are  $2 + 2n_1$ and  $2 + 2n'_1$ , respectively. Then  $n_1 = n'_1$ .

Because  $n_1 + n_2 + n_3 + n_4 = n'_1 + n'_2 + n'_3 + n'_4$ , by (15) and (16), we can deduce the following equations (17) and (18):

$$n_2n_3n_4 + (n_1 - 1 + q)(n_2n_3 + n_2n_4 + n_3n_4)$$

Luhua WANG and Ligong WANG

$$= n_2' n_3' n_4' + (n_1 - 1 + q) (n_2' n_3' + n_2' n_4' + n_3' n_4'),$$
(17)

$$[n_{1}(4-4(-1)^{q}-16q) + (2-2(-1)^{q}-16q)]n_{2}n_{3}n_{4} + [n_{1}(2-2(-1)^{q}-6q) + (1-(-1)^{q}-2q)](n_{2}n_{3}+n_{2}n_{4}+n_{3}n_{4}) = [n_{1}(4-4(-1)^{q}-16q) + (2-2(-1)^{q}-16q)]n_{2}'n_{3}'n_{4}' + [n_{1}(2-2(-1)^{q}-6q) + (1-(-1)^{q}-2q)] \cdot (n_{2}'n_{3}' + n_{2}'n_{4}' + n_{3}'n_{4}').$$
(18)

By solving the above two equations, we get that

$$\begin{cases} n_2 n_3 n_4 = n'_2 n'_3 n'_4, \\ n_2 n_3 + n_2 n_4 + n_3 n_4 = n'_2 n'_3 + n'_2 n'_4 + n'_3 n'_4. \end{cases}$$
(19)

Let

$$A = \prod_{i=2}^{4} n_i = \prod_{i=2}^{4} n'_i, \quad B = \sum_{2 \le i \le j \le 4} n_i n_j = \sum_{2 \le i \le j \le 4} n'_i n'_j.$$

Obviously,  $n_2, n_3, n_4$  and  $n'_2, n'_3, n'_4$  are the roots of the following equation, respectively,

 $x^{3} - (n - q - n_{1})x^{2} + Bx - A = 0.$ 

Hence the graph G' and G are isomorphic. Thus this theorem is proved.  $\Box$ 

**Theorem 3.4** Graph  $H(n; q, n_1, n_2, n_3, n_4)$  is determined by its Laplacian spectrum.

**Proof** Let  $G = H(n; q, n_1, n_2, n_3, n_4)$ . Assume that G' is Laplacian cospectral to G. By Lemma 2.12, G' is a connected unicyclic graph with n vertices, n edges and cycle  $C_q$ . Suppose that G' has  $x'_j$  vertices of degree j, for  $j = 1, 2, ..., \Delta$ , where  $\Delta$  is the maximum degree of G'. From Lemma 2.5, we obtain  $7 \leq \mu_1(G) = \mu_1(G') \leq 7 + \frac{1}{2}$ . Then  $\Delta = d_1(G') \leq 6$ . Because  $\max\{r_i(L_v(G))\} = 4$  for i = 1, 2, ..., n - 1, by Lemma 2.3, then  $\mu_1(L_v(G)) \leq 4$ . By Lemma 2.2, we can get  $\mu_2(L(G)) \leq \mu_1(L_v(G)) \leq \mu_1(L(G))$ , that is  $\mu_2(L(G)) \leq 4$ . Then according to Lemma 2.6,  $d_2(G') \leq \mu_2(L(G')) = \mu_2(L(G)) \leq 4$ . So, G' has at most one vertex of degree greater than 4. Therefore, by (i), (ii) and (v) of Lemma 2.1 and 2.12, we can get the following equations:

$$\begin{cases} x_1' + x_2' + x_3' + x_4' + 1 = n, \\ x_1' + 2x_2' + 3x_3' + 4x_4' + \Delta = 2n, \\ x_1' + 2^2x_2' + 3^2x_3' + 4^2x_4' + \Delta^2 = 4 + 6^2 + (n-5) \cdot 2^2, \\ x_1' + 2^3x_2' + 3^3x_3' + 4^3x_4' + \Delta^3 = 4 + 6^3 + (n-5) \cdot 2^3. \end{cases}$$
(20)

By solving Eqs. (20), we get the following result.

$$\begin{cases} x_1' = \frac{1}{6}\Delta^3 - \frac{3}{2}\Delta^2 + \frac{13}{3}\Delta - 4, \\ x_2' = n - \frac{1}{2}\Delta^3 + 4\Delta^2 - \frac{19}{2}\Delta + 16, \\ x_3' = \frac{1}{2}\Delta^3 - \frac{7}{2}\Delta^2 + 7\Delta - 24, \\ x_4' = 11 - \frac{1}{6}\Delta^3 + \Delta^2 - \frac{11}{6}\Delta. \end{cases}$$
(21)

Now we consider the following cases.

**Case 1** If  $\Delta = 1$ , then  $x'_1 = -1, x'_2 = n + 10, x'_3 = -20, x'_4 = 10.$ 

Laplacian spectral characterization of a kind of unicyclic graphs

**Case 2** If  $\Delta = 2$ , then  $x'_1 = 0, x'_2 = n + 9, x'_3 = -20, x'_4 = 10$ .

**Case 3** If  $\Delta = 3$ , then  $x'_1 = 0, x'_2 = n + 10, x'_3 = -21, x'_4 = 10$ .

**Case 4** If  $\Delta = 4$ , then  $x'_1 = 0, x'_2 = n + 10, x'_3 = -20, x'_4 = 9$ .

**Case 5** If  $\Delta = 6$ , then  $x'_1 = 4, x'_2 = n - 5, x'_3 = 0, x'_4 = 0$ .

Because  $x'_1$ ,  $x'_2$ ,  $x'_3$  and  $x'_4$  are nonnegative integers, we have  $\Delta = 6$ . Therefore, G' is only the graph  $H(n;q',n'_1,n'_2,n'_3,n'_4)$ . According to Theorem 3.3, we can know that the graph  $G' = H(n;q',n'_1,n'_2,n'_3,n'_4)$  is isomorphic to the graph  $G = H(n;q,n_1,n_2,n_3,n_4)$ . Thus, the theorem is proved.  $\Box$ 

For a graph, its Laplacian eigenvalues determine the eigenvalues of its complement [11], so the complements of all unicyclic graphs  $H(n; q, n_1, n_2, n_3, n_4)$  are determined by their Laplacian spectra.

Acknowledgement The authors would like to express their sincere gratitude to the referee for his/her careful reading of a previous version of this paper and for a number of helpful suggestions that led to many improvements.

### References

- G. AALIPOUR, S. AKBARI, N. SHAJARI. Laplacian spectral characterization of two families of trees. Linear Multilinear Algebra, 2013. Available from: <a href="http://dx.doi.org/10.1080/03081087.2013.801966">http://dx.doi.org/10.1080/03081087.2013.801966</a>>.
- [2] N. L. BIGGS. Algebraic Graph Theory. Combridge University Press, Cambridge, 1993.
- [3] A. E. BROUWER, W. H. HAEMERS. Spectra of Graphs. Springer, Berlin, 2012.
- [4] D. M. CVETKOVIĆ, A. R. BRUALDI. A Combinatorial Approach to Matrix Theory and its Application. CRC Press, Boca Raton, FL, 2009.
- [5] D. M. CVETKOVIĆ, M. DOOB, H. SACHS. Spectra of Graphs-Theory and Applications. III revised and enlarged edition, Johan Ambrosius Bart Verlag, Heidelberg-Leipzig, 1995.
- [6] E. R. van DAM, W. H. HAEMERS. Which graphs are determined by their spectrum? Linear Algebra Appl., 2003, 373: 241–272.
- [7] E. R. van DAM, W. H. HAEMERS. Developments on spectral characterizations of graphs. Discrete Math., 2009, 309: 576–586.
- [8] Jiming GUO. A conjecture on the algebraic connectivity of connected graphs with fixed girth. Discrete Math., 2008, 308(23): 5702–5711.
- [9] Jiming GUO. On the second largest Laplacian eigenvalue of trees. Linear Algebra Appl., 2005, 404: 251–261.
- [10] W. H. HAEMERS, Xiaogang LIU, Yuanping ZHANG. Spectral characterizations of lollipop graphs. Linear Algebra Appl., 2008, 428(11-12): 2415–2423.
- [11] A. K. KELMENS. The number of trees in a graph I. Automat. Remote Control, 1965, 26: 2118–2129.
- [12] A. K. KELMENS, V. M. CHELNOKOV. A certain polynomial of a graph and graphs with an extremal number of trees. J. Combinatorial Theory Ser. B, 1974, 16: 197–214.
- [13] Jiongsheng LI, Yongliang PAN. A note on the second largest eigenvalue of the Laplacian matrix of a graph. Linear Multilinear Algebra, 2000, 48(2): 117–121.
- [14] Jiongsheng LI, Xiaodong ZHANG. On the Laplacian eigenvalues of a graph. Linear Algebra Appl., 1998, 285(1-3): 305–307.
- [15] Fenjin LIU, Qiongxiang HUANG. On the spectral characterization of Π-shape trees. Linear Multilinear Algebra, 2013, 61(3): 355–367.
- [16] Fenjin LIU, Qiongxiang HUANG, Qinghai LIU. Spectral characterization of †-shape trees. Electron. J. Linear Algebra, 2011, 22: 822–837.
- [17] Xiaogang LIU, Suijie WANG. Laplacian spectral characterization of some graph products. Linear Algebra Appl., 2012, 437(7): 1749–1759.
- [18] Xiaogang LIU, Suijie WANG, Yuanping ZHANG. On the spectral characterization of some unicyclic graphs. Discrete Math., 2011, 311(21): 2317–2336.

- [19] Xiaogang LIU, Yuanping ZHANG, Pengli LU. One special double starlike graph is determined by its Laplacian spectrum. Appl. Math. Lett., 2009, 22(4): 435–438.
- [20] Pengli LU, Xuzhu WANG, Zuohan CHEN. Characterizing some class of unicyclic graph by its Laplacian spectrum. J. Harbin Engineering University, 2012, 33(7): 851–854.
- [21] C. S. OLIVEIRA, N. M. M. de ABREU, S. JURKIEWILZ. The characteristic polynomial of the Laplacian of graphs in (a, b)-linear cases. Linear Algebra Appl., 2012, 365: 113–121.
- [22] G. R. OMIDI, K. TAJBAKHSH. Starlike trees are determined by their Laplacian spectrum. Linear Algebra Appl., 2007, 422(2-3): 654–658.
- [23] Xiaoling SHEN, Yaoping HOU. A class of unicyclic graphs determined by their Laplacian spectrum. Electron. J. Linear Algebra, 2012, 23: 375–386.
- [24] Jianfeng WANG, Qiongxiang HUANG, F. BELARDA, et al. Spectral characterizations of dumbbell graphs. Electron. J. Combin., 2010, 17(1): Research Paper 42, 16 pp.
- [25] Jianfeng WANG, S. K. SIMIĆ, Qiongxiang HUANG, et al. Laplacian spectral characterization of disjoint union of paths and cycles. Linear Multilinear Algebra, 2011, 59(5): 531–539.
- [26] Wei WANG, Chengxian XU. On the spectral characterization of T-shape trees. Linear Algebra Appl., 2006, 414(2-3): 492–501.
- [27] Jiang ZHOU, Changjiang BU. Spectral characterization of line graphs of starlike trees. Linear Multilinear Algebra, 2013, 61(8): 1041–1050.
- [28] Jiang ZHOU, Changjiang BU. Laplacian spectral characterization of some graphs obtained by product operation. Discrete Math., 2012, 312(10): 1591–1595.