

Fekete-Szegő Problems for Certain Classes of Meromorphic Functions Using q -Derivative Operator

Huo TANG^{1,*}, H. M. ZAYED², A. O. MOSTAFA³, M. K. AOUF³

1. School of Mathematics and Statistics, Chifeng University, Inner Mongolia 024000, P. R. China;

2. Department of Mathematics, Faculty of Science, Menofia University, Shebin Elkom 32511, Egypt;

3. Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt

Abstract In this paper, we introduce two subclasses $\Sigma_q^*(\varphi)$ and $\Sigma_{q,\alpha}^*(\varphi)$ of meromorphic functions $f(z)$ for which $\frac{qzD_q f(z)}{f(z)} \prec \varphi(z)$ and

$$\frac{-(1 - \frac{\alpha}{q})qzD_q f(z) + \alpha qzD_q [zD_q f(z)]}{(1 - \frac{\alpha}{q})f(z) - \alpha zD_q f(z)} \prec \varphi(z), \quad \alpha \in \mathbb{C} \setminus (0, 1], \quad 0 < q < 1,$$

respectively. Sharp bounds for the Fekete-Szegő functional $|a_1 - \mu a_0^2|$ of the above classes are obtained. Also, we consider some applications of the results obtained to functions defined by q -Bessel function.

Keywords analytic function; meromorphic function; Fekete-Szegő problem; q -derivative operator; q -Bessel function

MR(2010) Subject Classification 30C45; 30C50

1. Introduction

The theory of q -analysis has important role in many areas of mathematics and physics, for example, in the areas of ordinary fractional calculus, optimal control problems, q -difference, q -integral equations and in q -transform analysis [1–4].

Let Σ denote the class of meromorphic functions of the form:

$$f(z) = \frac{1}{z} + \sum_{k=0}^{\infty} a_k z^k, \quad (1.1)$$

which are analytic in the open punctured unit disc $\mathbb{U}^* = \{z : z \in \mathbb{C} \text{ and } 0 < |z| < 1\} = \mathbb{U} \setminus \{0\}$.

A function $f \in \Sigma$ is meromorphic starlike of order α , denoted by $\Sigma^*(\alpha)$, if

$$-\Re\left\{\frac{zf'(z)}{f(z)}\right\} > \alpha, \quad 0 \leq \alpha < 1; \quad z \in \mathbb{U}.$$

Received April 26, 2017; Accepted March 23, 2018

Supported by the National Natural Science Foundation of China (Grant Nos. 11561001; 11271045), the Program for Young Talents of Science and Technology in Universities of Inner Mongolia Autonomous Region (Grant No. NJYT-18-A14), the Natural Science Foundation of Inner Mongolia (Grant Nos. 2010MS0117; 2014MS0101; 2017MS0113), the Higher School Foundation of Inner Mongolia (Grant Nos. NJZY16251; NJZY17300; NJZY17301; NJZY18217) and the Natural Science Foundation of Chifeng.

* Corresponding author

E-mail address: thth2009@163.com (Huo TANG); hanaa.zayed42@yahoo.com (H. M. ZAYED); adelaeg254@yahoo.com (A. O. MOSTAFA); mkaouf127@yahoo.com (M. K. AOUF)

The class $\Sigma^*(\alpha)$ was introduced and studied by Pommerenke [5] (see also Miller [6]).

Let $\varphi(z)$ be an analytic function with positive real part on \mathbb{U} satisfying $\varphi(0) = 1$ and $\varphi'(0) > 0$ which maps \mathbb{U} onto a region starlike with respect to 1 and symmetric with respect to the real axis.

Let $\Sigma^*(\varphi)$ be the class of functions $f \in \Sigma$ for which

$$-\frac{zf'(z)}{f(z)} \prec \varphi(z), \quad z \in \mathbb{U}.$$

The class $\Sigma^*(\varphi)$ was introduced and studied by Silverman et al. [7]. We note that, the class $\Sigma^*(\alpha)$ is the special case of the class $\Sigma^*(\varphi)$ when $\varphi(z) = \frac{1+(1-2\alpha)z}{1-z}$ ($0 \leq \alpha < 1$).

For a function $f(z) \in \Sigma$ given by (1.1) and $0 < q < 1$, the q -derivative of a function $f(z)$ is defined by [2]

$$D_q f(z) = \frac{f(qz) - f(z)}{(q-1)z}, \quad z \in \mathbb{U}^*. \tag{1.2}$$

From (1.2), we deduce that $D_q f(z)$ for a function $f(z)$ of the form (1.1) is given by

$$D_q f(z) = -\frac{1}{qz^2} + \sum_{k=0}^{\infty} [k]_q a_k z^{k-1}, \quad z \neq 0,$$

where

$$[k]_q = \frac{1 - q^k}{1 - q}.$$

As $q \rightarrow 1^-$, $[k]_q \rightarrow k$, we have $\lim_{q \rightarrow 1^-} D_q f(z) = f'(z)$.

Making use of the q -derivative D_q , we introduce the subclasses $\Sigma_q^*(\varphi)$ and $\Sigma_{q,\alpha}^*(\varphi)$ as follows:

Definition 1.1 A function $f(z) \in \Sigma$ is said to be in the class $\Sigma_q^*(\varphi)$, if and only if

$$-\frac{qzD_q f(z)}{f(z)} \prec \varphi(z), \quad z \in \mathbb{U}. \tag{1.3}$$

We note that:

- (i) $\lim_{q \rightarrow 1^-} \Sigma_q^*(\varphi) = \Sigma^*(\varphi)$ (see [7] and [8, with $\alpha = 0$]);
- (ii) $\lim_{q \rightarrow 1^-} \Sigma_q^*\left(\frac{1+(1-2\alpha)z}{1-z}\right) = \Sigma^*(\alpha)$ ($0 \leq \alpha < 1$) (see [5]);
- (iii) $\lim_{q \rightarrow 1^-} \Sigma_q^*\left(\frac{1+z}{1-z}\right) = \mathcal{F}^*(1) = \mathcal{F}^*$ (see [9, with $b = 1$]);
- (iv) $\lim_{q \rightarrow 1^-} \Sigma_q^*\left(\frac{1+\beta(1-2\gamma\delta)z}{1+\beta(1-2\gamma)z}\right) = \Sigma(\delta, \beta, \gamma)$ ($0 \leq \delta < 1$, $0 < \beta \leq 1$, $\frac{1}{2} \leq \gamma \leq 1$) (see [10]);
- (v) $\lim_{q \rightarrow 1^-} \Sigma_q^*\left(\frac{1+Az}{1+Bz}\right) = K_1(A, B)$ ($0 \leq B < 1$, $-B < A < B$) (see [11]).

Definition 1.2 For $\alpha \in \mathbb{C} \setminus (0, 1]$, let $\Sigma_{q,\alpha}^*(\varphi)$ be the subclass of Σ consisting of functions $f(z)$ of the form (1.1) and satisfying the analytic criterion:

$$\frac{-(1 - \frac{\alpha}{q})qzD_q f(z) + \alpha qzD_q[zD_q f(z)]}{(1 - \frac{\alpha}{q})f(z) - \alpha zD_q f(z)} \prec \varphi(z). \tag{1.4}$$

From (1.3) and (1.4), we note that

$$\Sigma_{q,0}^*(\varphi) = \Sigma_q^*(\varphi)$$

and $\lim_{q \rightarrow 1^-} \Sigma_{q,\alpha}^*(\varphi) = \Sigma_\alpha^*(\varphi)$ (see [7]).

2. Fekete-Szegő problems

To prove our results, we need the following lemmas.

Lemma 2.1 ([12]) *If $p(z) = 1 + c_1z + c_2z^2 + \dots$ is a function with positive real part in \mathbb{U} and μ is a complex number, then*

$$|c_2 - \mu c_1^2| \leq 2 \max\{1; |2\mu - 1|\}.$$

The result is sharp for the functions given by

$$p(z) = \frac{1 + z^2}{1 - z^2} \text{ and } p(z) = \frac{1 + z}{1 - z}.$$

Lemma 2.2 ([12]) *If $p_1(z) = 1 + c_1z + c_2z^2 + \dots$ is a function with positive real part in \mathbb{U} , then*

$$|c_2 - \nu c_1^2| \leq \begin{cases} -4\nu + 2, & \text{if } \nu \leq 0, \\ 2, & \text{if } 0 \leq \nu \leq 1, \\ 4\nu - 2, & \text{if } \nu \geq 1. \end{cases}$$

When $\nu < 0$ or $\nu > 1$, the equality holds if and only if $p_1(z) = \frac{1+z}{1-z}$ or one of its rotations. If $0 < \nu < 1$, then the equality holds if and only if $p_1(z) = \frac{1+z^2}{1-z^2}$ or one of its rotations. If $\nu = 0$, the equality holds if and only if

$$p_1(z) = \left(\frac{1}{2} + \frac{\lambda}{2}\right) \frac{1+z}{1-z} + \left(\frac{1}{2} - \frac{\lambda}{2}\right) \frac{1-z}{1+z}, \quad 0 \leq \lambda \leq 1,$$

or one of its rotations. If $\nu = 1$, the equality holds if and only if

$$\frac{1}{p_1(z)} = \left(\frac{1}{2} + \frac{\lambda}{2}\right) \frac{1+z}{1-z} + \left(\frac{1}{2} - \frac{\lambda}{2}\right) \frac{1-z}{1+z}, \quad 0 \leq \lambda \leq 1,$$

or one of its rotations. Also the above upper bound is sharp and it can be improved as follows when $0 < \nu < 1$:

$$|c_2 - \nu c_1^2| + \nu |c_1|^2 \leq 2, \quad 0 < \nu \leq \frac{1}{2}$$

and

$$|c_2 - \nu c_1^2| + (1 - \nu) |c_1|^2 \leq 2, \quad \frac{1}{2} < \nu < 1.$$

Unless otherwise mentioned, we assume throughout this paper that $\alpha \in \mathbb{C} \setminus (0, 1]$ and $0 < q < 1$.

Theorem 2.3 *Let $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$ ($B_1 \geq 0$). If $f(z)$ given by (1.1) belongs to the class $\Sigma_q^*(\varphi)$ and μ is a complex number, then*

$$|a_1 - \mu a_0^2| \leq \frac{|B_1|}{(1+q)} \max\{1, \left|\frac{B_2}{B_1} - [1 - \mu(1+q)]B_1\right|\}, \quad B_1 \neq 0, \tag{2.1}$$

$$|a_1| \leq \frac{|B_2|}{(1+q)}, \quad B_1 = 0. \tag{2.2}$$

The result is sharp.

Proof If $f(z) \in \Sigma_q^*(\varphi)$, then there is a Schwarz function $w(z)$ in \mathbb{U} with $w(0) = 0$ and $|w(z)| < 1$ in \mathbb{U} and such that

$$-\frac{qzD_q f(z)}{f(z)} = \varphi(w(z)). \tag{2.3}$$

Define the function $p_1(z)$ by

$$p_1(z) = \frac{1 + w(z)}{1 - w(z)} = 1 + c_1z + c_2z^2 + \dots \tag{2.4}$$

Since $w(z)$ is a Schwarz function, we see that $\Re\{p_1(z)\} > 0$ and $p_1(0) = 1$.

Define

$$p(z) = -\frac{qzD_q f(z)}{f(z)} = 1 + b_1z + b_2z^2 + \dots \tag{2.5}$$

In view of (2.3)–(2.5), we have

$$p(z) = \varphi\left(\frac{p_1(z) - 1}{p_1(z) + 1}\right). \tag{2.6}$$

Since

$$\frac{p_1(z) - 1}{p_1(z) + 1} = \frac{1}{2}[c_1z + (c_2 - \frac{c_1^2}{2})z^2 + (c_3 + \frac{c_1^3}{4} - c_1c_2)z^3 + \dots].$$

Therefore, we have

$$\varphi\left(\frac{p_1(z) - 1}{p_1(z) + 1}\right) = 1 + \frac{1}{2}B_1c_1z + \left[\frac{1}{2}B_1(c_2 - \frac{c_1^2}{2}) + \frac{1}{4}B_2c_1^2\right]z^2 + \dots \tag{2.7}$$

From (2.6) and (2.7), we obtain

$$b_1 = \frac{1}{2}B_1c_1,$$

and

$$b_2 = \frac{1}{2}B_1(c_2 - \frac{c_1^2}{2}) + \frac{1}{4}B_2c_1^2.$$

Then, from (2.5) and (1.1), we see that $b_1 = -a_0$, and $b_2 = a_0^2 - (q + 1)a_1$, or, equivalently, we have

$$a_0 = -\frac{B_1c_1}{2}, \tag{2.8}$$

and

$$a_1 = -\frac{B_1}{2(1+q)}\left[c_2 - \frac{c_1^2}{2}\left(1 - \frac{B_2}{B_1} + B_1\right)\right]. \tag{2.9}$$

Therefore

$$a_1 - \mu a_0^2 = -\frac{B_1}{2(1+q)}\{c_2 - \nu c_1^2\},$$

where

$$\nu = \frac{1}{2}\left[1 - \frac{B_2}{B_1} + B_1 - \mu B_1(1+q)\right]. \tag{2.10}$$

Now, the result (2.1) follows by an application of Lemma 2.1. Also, if $B_1 = 0$, then

$$a_0 = 0, \quad a_1 = -\frac{B_2c_1^2}{4(1+q)}.$$

Since $p(z)$ has positive real part, then $|c_1| \leq 2$ (see [13]). Hence

$$|a_1| \leq \frac{|B_2|}{1+q},$$

this proving (2.2). The result is sharp for the functions

$$-\frac{qzD_q f(z)}{f(z)} = \varphi(z^2), \quad -\frac{qzD_q f(z)}{f(z)} = \varphi(z).$$

This completes the proof of Theorem 2.3. \square

Remark 2.4 For $q \rightarrow 1^-$ in Theorem 2.3, we obtain the result obtained by [7, Theorem 2.1].

Putting $q \rightarrow 1^-$ and $\varphi(z) = \frac{1+z}{1-z}$ in Theorem 2.3, we obtain the following corollary.

Corollary 2.5 If $f(z)$ given by (1.1) belongs to the class \mathcal{F}^* and μ is a complex number, then

$$|a_1 - \mu a_0^2| \leq \max\{1, |1 - 2(1 - 2\mu)|\}.$$

The result is sharp.

By using Lemma 2.2, we can obtain the following theorem.

Theorem 2.6 Let $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$ ($B_i > 0, i = 1, 2$). If $f(z)$ given by (1.1) belongs to the class $\Sigma_q^*(\varphi)$, then

$$|a_1 - \mu a_0^2| \leq \begin{cases} \frac{B_2 - [1 - \mu(1+q)]B_1^2}{1+q}, & \text{if } \mu \leq \sigma_1, \\ \frac{B_1}{1+q}, & \text{if } \sigma_1 \leq \mu \leq \sigma_2, \\ \frac{-B_2 + [1 - \mu(1+q)]B_1^2}{1+q}, & \text{if } \mu \geq \sigma_2, \end{cases} \quad (2.11)$$

where

$$\sigma_1 = \frac{-B_1 - B_2 + B_1^2}{(1+q)B_1^2}, \quad \sigma_2 = \frac{B_1 - B_2 + B_1^2}{(1+q)B_1^2}.$$

The result is sharp. Further, let $\sigma_3 = \frac{-B_2 + B_1^2}{(1+q)B_1^2}$.

(i) If $\sigma_1 \leq \mu \leq \sigma_3$, then

$$|a_1 - \mu a_0^2| + \frac{\{(B_1 + B_2) + [\mu(q + 1) - 1]B_1^2\}|a_0|^2}{(1+q)B_1^2} \leq \frac{B_1}{1+q}. \quad (2.12)$$

(ii) If $\sigma_3 \leq \mu \leq \sigma_2$, then

$$|a_1 - \mu a_0^2| + \frac{\{(B_1 - B_2) + [1 - \mu(q + 1)]B_1^2\}|a_0|^2}{(1+q)B_1^2} \leq \frac{B_1}{1+q}. \quad (2.13)$$

Proof First, let $\mu \leq \sigma_1$. Then

$$\begin{aligned} |a_1 - \mu a_0^2| &\leq \frac{B_1}{(1+q)} \left\{ \frac{B_2}{B_1} - [1 - \mu(1+q)]B_1 \right\} \\ &\leq \frac{B_2 - [1 - \mu(1+q)]B_1^2}{1+q}. \end{aligned}$$

Let, now $\sigma_1 \leq \mu \leq \sigma_2$. Then, using the above calculations, we obtain

$$|a_1 - \mu a_0^2| \leq \frac{B_1}{1+q}.$$

Finally, if $\mu \geq \sigma_2$, then

$$|a_1 - \mu a_0^2| \leq \frac{B_1}{(1+q)} \left\{ -\frac{B_2}{B_1} + [1 - \mu(1+q)]B_1 \right\} \leq \frac{-B_2 + [1 - \mu(1+q)]B_1^2}{1+q}.$$

To show that the bounds are sharp, we define the functions $K_{\varphi n}$ ($n \geq 2$) by

$$-\frac{qzD_qK_{\varphi n}(z)}{K_{\varphi n}(z)} = \varphi(z^{n-1})(z^2K_{\varphi n}(z)|_{z=0} = 0 = -z^2K'_{\varphi n}(z)|_{z=0} - 1)$$

and the functions F_γ and G_γ ($0 \leq \gamma \leq 1$) by

$$-\frac{qzD_qF_\gamma(z)}{F_\gamma(z)} = \varphi\left(\frac{z(z+\gamma)}{1+\gamma z}\right)(z^2F_\gamma(z)|_{z=0} = 0 = -z^2F'_\gamma(z)|_{z=0} - 1)$$

and

$$-\frac{qzD_qG_\gamma(z)}{G_\gamma(z)} = \varphi\left(-\frac{z(z+\gamma)}{1+\gamma z}\right)(z^2G_\gamma(z)|_{z=0} = 0 = -z^2G'_\gamma(z)|_{z=0} - 1).$$

Clearly, the functions $K_{\varphi n}$, F_γ and $G_\gamma \in \Sigma_q^*(\varphi)$. Also we write $K_\varphi = K_{\varphi 2}$. If $\mu < \sigma_1$ or $\mu > \sigma_2$, then the equality holds if and only if $f(z)$ is K_φ or one of its rotations. When $\sigma_1 < \mu < \sigma_2$, then the equality holds if $f(z)$ is $K_{\varphi 3}$ or one of its rotations. If $\mu = \sigma_1$, then the equality holds if and only if $f(z)$ is F_γ or one of its rotations. If $\mu = \sigma_2$, then the equality holds if and only if $f(z)$ is G_γ or one of its rotations. This completes the proof of Theorem 2.6. \square

Remark 2.7 (i) For $q \rightarrow 1^-$ in Theorem 2.6, we obtain the result obtained by [8, Theorem 5.1];

(ii) Putting $q \rightarrow 1^-$ and $\varphi(z) = \frac{1+z}{1-z}$ in Theorem 2.6, we obtain a new result for the class \mathcal{F}^* .

Theorem 2.8 Let $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$ ($B_1 \geq 0$). If $f(z)$ given by (1.1) belongs to the class $\Sigma_{q,\alpha}^*(\varphi)$ and μ is a complex number, then

$$|a_1 - \mu a_0^2| \leq \left(\frac{q}{1+q}\right) \left|\frac{B_1}{q - \alpha(1+q)}\right| \max\left\{1, \left|\frac{B_2}{B_1} - \left[1 - \mu \frac{q(1+q)[q - \alpha(1+q)]}{(q - \alpha)^2}\right] B_1\right|\right\}, \quad B_1 \neq 0, \tag{2.14}$$

$$|a_1| \leq \left(\frac{q}{1+q}\right) \left|\frac{B_2}{q - \alpha(1+q)}\right|, \quad B_1 = 0. \tag{2.15}$$

The result is sharp.

Proof If $f(z) \in \Sigma_{q,\alpha}^*(\varphi)$, then there is a Schwarz function $w(z)$ in \mathbb{U} with $w(0) = 0$ and $|w(z)| < 1$ in \mathbb{U} and such that

$$\frac{-(1 - \frac{\alpha}{q})qzD_qf(z) + \alpha qzD_q[zD_qf(z)]}{(1 - \frac{\alpha}{q})f(z) - \alpha zD_qf(z)} = \varphi(w(z)).$$

Since $w(z)$ is a Schwarz function, we see that $\Re\{p_1(z)\} > 0$ and $p_1(0) = 1$.

Define

$$p(z) = \frac{-(1 - \frac{\alpha}{q})qzD_qf(z) + \alpha qzD_q[zD_qf(z)]}{(1 - \frac{\alpha}{q})f(z) - \alpha zD_qf(z)} = 1 + b_1z + b_2z^2 + \dots \tag{2.16}$$

Then, from (2.6), (2.7), (2.16) and (1.1), we see that

$$\frac{1}{2}B_1c_1 = -(1 - \frac{\alpha}{q})a_0,$$

and

$$\frac{1}{2}B_1(c_2 - \frac{c_1^2}{2}) + \frac{1}{4}B_2c_1^2 = (1 - \frac{\alpha}{q})^2a_0^2 - (1+q)(1 - \alpha - \frac{\alpha}{q})a_1,$$

or, equivalently, we have

$$a_0 = -\frac{qB_1c_1}{2(q-\alpha)}, \quad a_1 = -\frac{qB_1}{2(1+q)[q-\alpha(1+q)]} \left[c_2 - \frac{c_1^2}{2} \left(1 - \frac{B_2}{B_1} + B_1 \right) \right].$$

Therefore

$$a_1 - \mu a_0^2 = -\frac{qB_1}{2(1+q)[q-\alpha(1+q)]} \{ c_2 - \nu c_1^2 \},$$

where

$$\nu = \frac{1}{2} \left[1 - \frac{B_2}{B_1} + B_1 \left(1 - \mu \frac{q(1+q)[q-\alpha(1+q)]}{(q-\alpha)^2} \right) \right]. \tag{2.17}$$

Now, the result (2.14) follows by an application of Lemma 2.1. Also, if $B_1 = 0$, then

$$a_0 = 0, \quad a_1 = \frac{qB_2c_1^2}{4(1+q)[q-\alpha(1+q)]}.$$

Since $p(z)$ has positive real part, then $|c_1| \leq 2$ (see [13]), hence

$$|a_1| \leq \left(\frac{q}{1+q} \right) \left| \frac{B_2}{q-\alpha(1+q)} \right|,$$

this proving (2.15). The result is sharp for the functions

$$\frac{-(1-\frac{\alpha}{q})qzD_qf(z) + \alpha qzD_q[zD_qf(z)]}{(1-\frac{\alpha}{q})f(z) - \alpha zD_qf(z)} = \varphi(z^2)$$

and

$$\frac{-(1-\frac{\alpha}{q})qzD_qf(z) + \alpha qzD_q[zD_qf(z)]}{(1-\frac{\alpha}{q})f(z) - \alpha zD_qf(z)} = \varphi(z).$$

This completes the proof of Theorem 2.8. \square

Remark 2.9 (i) Putting $\alpha = 0$ in Theorem 2.8, we obtain the result of Theorem 2.3;

(ii) Putting $q \rightarrow 1^-$ in Theorem 2.8, we obtain the result obtained by [7, Theorem 2.2].

Using arguments similar to those in the proof of Theorem 2.6, we obtain the following theorem.

Theorem 2.10 Let $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$ ($B_i > 0, i \in 1, 2, 0 < \alpha < \frac{q}{1+q}$). If $f(z)$ given by (1.1) belongs to the class $\Sigma_{q,\alpha}^*(\varphi)$, then

$$|a_1 - \mu a_0^2| \leq \begin{cases} \frac{q}{(1+q)[q-\alpha(1+q)]} \{ B_2 - [1 - \mu \frac{q(1+q)[q-\alpha(1+q)]}{(q-\alpha)^2}] B_1^2 \}, & \text{if } \mu \leq \sigma_4, \\ \frac{qB_1}{(1+q)[q-\alpha(1+q)]}, & \text{if } \sigma_4 \leq \mu \leq \sigma_5, \\ \frac{q}{(1+q)[q-\alpha(1+q)]} \{ -B_2 + [1 - \mu \frac{q(1+q)[q-\alpha(1+q)]}{(q-\alpha)^2}] B_1^2 \}, & \text{if } \mu \geq \sigma_5, \end{cases}$$

where

$$\sigma_4 = \frac{(q-\alpha)^2[-B_1 - B_2 + B_1^2]}{q(1+q)[q-\alpha(1+q)]B_1^2}, \quad \sigma_5 = \frac{(q-\alpha)^2[B_1 - B_2 + B_1^2]}{q(1+q)[q-\alpha(1+q)]B_1^2}.$$

The result is sharp. Further, let

$$\sigma_6 = \frac{(q-\alpha)^2[-B_2 + B_1^2]}{q(1+q)[q-\alpha(1+q)]B_1^2}.$$

(i) If $\sigma_4 \leq \mu \leq \sigma_6$, then

$$|a_1 - \mu a_0^2| + \frac{(q - \alpha)^2}{q(1 + q)[q - \alpha(1 + q)]B_1^2} \times \{(B_1 + B_2) + [\mu \frac{q(1 + q)[q - \alpha(1 + q)]}{(q - \alpha)^2} - 1]B_1^2\} |a_0|^2 \leq \frac{qB_1}{(1 + q)[q - \alpha(1 + q)]}.$$

(ii) If $\sigma_6 \leq \mu \leq \sigma_5$, then

$$|a_1 - \mu a_0^2| + \frac{(q - \alpha)^2}{q(1 + q)[q - \alpha(1 + q)]B_1^2} \times \{(B_1 - B_2) + [1 - \mu \frac{q(1 + q)[q - \alpha(1 + q)]}{(q - \alpha)^2}]B_1^2\} |a_0|^2 \leq \frac{qB_1}{(1 + q)[q - \alpha(1 + q)]}.$$

3. Applications to functions defined by q -Bessel function

We recall some definitions of q -calculus which will be used in our paper.

For any complex number α , the q -shifted factorials are defined by

$$(\alpha; q)_0 = 1; (\alpha; q)_n = \prod_{k=0}^{n-1} (1 - \alpha q^k), \quad n \in \mathbb{N} = \{1, 2, \dots\}. \tag{3.1}$$

If $|q| < 1$, the definition (3.1) remains meaningful for $n = \infty$ as a convergent infinite product

$$(\alpha; q)_\infty = \prod_{j=0}^{\infty} (1 - \alpha q^j).$$

In terms of the analogue of the gamma function

$$(q^\alpha; q)_n = \frac{\Gamma_q(\alpha + n)(1 - q)^n}{\Gamma_q(\alpha)}, \quad n > 0,$$

where the q -gamma function is defined by

$$\Gamma_q(x) = \frac{(q; q)_\infty (1 - q)^{1-x}}{(q^x; q)_\infty}, \quad 0 < q < 1.$$

We note that $\lim_{q \rightarrow 1^-} \frac{(q^\alpha; q)_n}{(1 - q)^n} = (\alpha)_n$, where

$$(\alpha)_n = \begin{cases} 1, & \text{if } n = 0, \\ \alpha(\alpha + 1)(\alpha + 2) \cdots (\alpha + n - 1), & \text{if } n \in \mathbb{N}. \end{cases}$$

Now, consider the q -analogue of Bessel function defined by [3]

$$\mathcal{J}_\nu^{(1)}(z; q) = \frac{(q^{\nu+1}; q)_\infty}{(q; q)_\infty} \sum_{k=0}^{\infty} \frac{(-1)^k}{(q; q)_k (q^{\nu+1}; q)_k} \left(\frac{z}{2}\right)^{2k+\nu}, \quad 0 < q < 1.$$

Also, let us define

$$\begin{aligned} \mathcal{L}_\nu(z; q) &= \frac{2^\nu (q; q)_\infty}{(q^{\nu+1}; q)_\infty (1 - q)^{\nu} z^{\nu/2+1}} \mathcal{J}_\nu^{(1)}(z^{1/2}(1 - q); q) \\ &= \frac{1}{z} + \sum_{k=0}^{\infty} \frac{(-1)^{k+1} (1 - q)^{2(k+1)}}{4^{(k+1)} (q; q)_{k+1} (q^{\nu+1}; q)_{k+1}} z^k, \quad z \in \mathbb{U}. \end{aligned}$$

By using the Hadamard product (or convolution), we define the linear operator $\mathcal{L}_{q,v} : \Sigma \rightarrow \Sigma$, as follows:

$$(\mathcal{L}_{q,v}f)(z) = \mathcal{L}_v(z; q) * f(z) = \frac{1}{z} + \sum_{k=0}^{\infty} \frac{(-1)^{k+1}(1-q)^{2(k+1)}}{4^{(k+1)}(q; q)_{k+1}(q^{v+1}; q)_{k+1}} a_k z^k.$$

The linear operator $\mathcal{L}_{q,v}$ was introduced and studied by Mostafa et al. [14]. Also, as $q \rightarrow 1^-$, the linear operator $\mathcal{L}_{q,v}$ reduces to the operator \mathcal{L}_v introduced and studied by Aouf et al. [15].

For $0 < q < 1$ and $\alpha \in \mathbb{C} \setminus (0, 1]$, let $\Sigma_{q,v}^*(\varphi)$ and $\Sigma_{q,\alpha,v}^*(\varphi)$ be the subclasses of Σ consisting of functions $f(z)$ of the form (1.1) and satisfy the analytic criterions, respectively:

$$-\frac{qzD_q(\mathcal{L}_{q,v}f)(z)}{(\mathcal{L}_{q,v}f)(z)} \prec \varphi(z), \quad z \in \mathbb{U},$$

$$\frac{-(1 - \frac{\alpha}{q})qzD_q(\mathcal{L}_{q,v}f) + \alpha qzD_q[zD_q(\mathcal{L}_{q,v}f)]}{(1 - \frac{\alpha}{q})f(z) - \alpha zD_q(\mathcal{L}_{q,v}f)(z)} \prec \varphi(z), \quad z \in \mathbb{U}.$$

Using similar arguments to those in the proof of the above theorems, we obtain the following theorems.

Theorem 3.1 *Let $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$ ($B_1 \geq 0$). If $f(z)$ given by (1.1) belongs to the class $\Sigma_{q,v}^*(\varphi)$ and μ is a complex number, then*

$$|a_1 - \mu a_0^2| \leq \frac{4^2(1 - q^{v+1})(1 - q^{v+2})|B_1|}{(1 - q)^2} \times$$

$$\max\{1, |\frac{B_2}{B_1} - [1 - \mu(\frac{1 - q^{v+1}}{1 - q^{v+2}})]B_1|\}, \quad B_1 \neq 0,$$

$$|a_1| \leq \frac{4^2(1 - q^{v+1})(1 - q^{v+2})|B_2|}{(1 - q)^2}, \quad B_1 = 0.$$

The result is sharp.

Theorem 3.2 *Let $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$ ($B_i > 0, i = 1, 2$). If $f(z)$ given by (1.1) belongs to the class $\Sigma_{q,v}^*(\varphi)$, then*

$$|a_1 - \mu a_0^2| \leq \begin{cases} \frac{4^2(1 - q^{v+1})(1 - q^{v+2})}{(1 - q)^2} \{B_2 - [1 - \mu(\frac{1 - q^{v+1}}{1 - q^{v+2}})]B_1^2\}, & \text{if } \mu \leq \sigma_1^*, \\ \frac{4^2(1 - q^{v+1})(1 - q^{v+2})B_1}{(1 - q)^2}, & \text{if } \sigma_1^* \leq \mu \leq \sigma_2^*, \\ \frac{4^2(1 - q^{v+1})(1 - q^{v+2})}{(1 - q)^2} \{-B_2 + [1 - \mu(\frac{1 - q^{v+1}}{1 - q^{v+2}})]B_1^2\}, & \text{if } \mu \geq \sigma_2^*, \end{cases}$$

where

$$\sigma_1^* = \frac{[-B_1 - B_2 + B_1^2](1 - q^{v+2})}{(1 - q^{v+1})B_1^2}, \quad \sigma_2^* = \frac{[B_1 - B_2 + B_1^2](1 - q^{v+2})}{(1 - q^{v+1})B_1^2}.$$

The result is sharp. Further, let

$$\sigma_3^* = \frac{[-B_2 + B_1^2](1 - q^{v+2})}{(1 - q^{v+1})B_1^2}.$$

(i) If $\sigma_1^* \leq \mu \leq \sigma_3^*$, then

$$|a_1 - \mu a_0^2| + \frac{(1 - q^{v+2})}{(1 - q^{v+1})B_1^2} \{ (B_1 + B_2) + [\mu(\frac{1 - q^{v+1}}{1 - q^{v+2}}) - 1]B_1^2 \} |a_0|^2 \leq \frac{4^2(1 - q^{v+1})(1 - q^{v+2})B_1}{(1 - q)^2}.$$

(ii) If $\sigma_3^* \leq \mu \leq \sigma_2^*$, then

$$|a_1 - \mu a_0^2| + \frac{(1 - q^{v+2})}{(1 - q^{v+1})B_1^2} \{ (B_1 - B_2) + [1 - \mu(\frac{1 - q^{v+1}}{1 - q^{v+2}})]B_1^2 \} |a_0|^2 \leq \frac{4^2(1 - q^{v+1})(1 - q^{v+2})B_1}{(1 - q)^2}.$$

Theorem 3.3 Let $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$ ($B_1 \geq 0$). If $f(z)$ given by (1.1) belongs to the class $\Sigma_{q,\alpha,v}^*(\varphi)$ and μ is a complex number, then

$$|a_1 - \mu a_0^2| \leq \frac{4^2q(1 - q^{v+1})(1 - q^{v+2})|B_1|}{(1 - q)^2|q - \alpha(1 + q)|} \times \max\{1, |\frac{B_2}{B_1} - [1 - \mu \frac{q(1 - q^{v+1})[q - \alpha(1 + q)]}{(\alpha - q)^2(1 - q^{v+2})}]B_1|\}, \quad B_1 \neq 0,$$

$$|a_1| \leq \frac{4^2q(1 - q^{v+1})(1 - q^{v+2})|B_1|}{(1 - q)^2|q - \alpha(1 + q)|}, \quad B_1 = 0.$$

The result is sharp.

Theorem 3.4 Let $\varphi(z) = 1 + B_1z + B_2z^2 + \dots$ ($B_i > 0, i = 1, 2, 0 < \alpha < \frac{q}{1+q}$). If $f(z)$ given by (1.1) belongs to the class $\Sigma_{q,\alpha,v}^*(\varphi)$, then

$$|a_1 - \mu a_0^2| \leq \begin{cases} \frac{4^2(1 - q^{v+1})(1 - q^{v+2})}{(1 - q)^2|q - \alpha(1 + q)|} \times \{ B_2 - [1 - \mu \frac{[\alpha + (1 - \alpha)q](1 - 2\alpha)(1 - q^{v+1})}{q(1 - \alpha)^2(1 - q^{v+2})}]B_1^2 \}, & \text{if } \mu \leq \sigma_4^*, \\ \frac{4^2(1 - q^{v+1})(1 - q^{v+2})B_1}{(1 - q)^2|q - \alpha(1 + q)|}, & \text{if } \sigma_4^* \leq \mu \leq \sigma_5^*, \\ \frac{4^2(1 - q^{v+1})(1 - q^{v+2})}{(1 - q)^2|q - \alpha(1 + q)|} \times \{ -B_2 + [1 - \mu \frac{q(1 - q^{v+1})[q - \alpha(1 + q)]}{(q - \alpha)^2(1 - q^{v+2})}]B_1^2 \}, & \text{if } \mu \geq \sigma_5^*, \end{cases}$$

where

$$\sigma_4^* = \frac{(q - \alpha)^2(1 - q^{v+2})[-B_1 - B_2 + B_1^2]}{qB_1^2(1 - q^{v+1})[q - \alpha(1 + q)]},$$

$$\sigma_5^* = \frac{(q - \alpha)^2(1 - q^{v+2})[B_1 - B_2 + B_1^2]}{qB_1^2(1 - q^{v+1})[q - \alpha(1 + q)]}.$$

The result is sharp. Further, let

$$\sigma_6^* = \frac{(q - \alpha)^2(1 - q^{v+2})[-B_2 + B_1^2]}{qB_1^2(1 - q^{v+1})[q - \alpha(1 + q)]}.$$

(i) If $\sigma_4^* \leq \mu \leq \sigma_6^*$, then

$$\begin{aligned} & |a_1 - \mu a_0^2| + \frac{(q - \alpha)^2(1 - q^{v+2})}{q(1 - q^{v+1})[q - \alpha(1 + q)]B_1^2} \times \\ & \quad \{(B_1 + B_2) + [\mu \frac{q(1 - q^{v+2})[q - \alpha(1 + q)]}{(1 - q^{v+1})(\alpha - q)^2} - 1]B_1^2\} |a_0|^2 \\ & \leq \frac{4^2(1 - q^{v+1})(1 - q^{v+2})B_1}{(1 - q)^2[q - \alpha(1 + q)]}. \end{aligned}$$

(ii) If $\sigma_6^* \leq \mu \leq \sigma_5^*$, then

$$\begin{aligned} & |a_1 - \mu a_0^2| + \frac{(q - \alpha)^2(1 - q^{v+2})}{q(1 - q^{v+1})[q - \alpha(1 + q)]B_1^2} \times \\ & \quad \{(B_1 - B_2) + [1 - \mu \frac{q(1 - q^{v+2})[q - \alpha(1 + q)]}{(1 - q^{v+1})(\alpha - q)^2}]B_1^2\} |a_0|^2 \\ & \leq \frac{4^2(1 - q^{v+1})(1 - q^{v+2})B_1}{(1 - q)^2[q - \alpha(1 + q)]}. \end{aligned}$$

Acknowledgements We would like to thank the referees for their valuable comments, suggestions and corrections.

References

- [1] M. H. ABU-RISHA, M. H. ANNABY, M. E. H. ISMAIL, et al. *Linear q-difference equations*. Z. Anal. Anwend., 2007, **26**(4): 481–494.
- [2] G. GASPER, M. RAHMAN. *Basic Hypergeometric Series*. Cambridge University Press, Cambridge, 1990.
- [3] F. H. JACKSON. *The application of basic numbers to Bessel's and Legendre's functions*. Proc. London Math. Soc. (2), 1905, **3**: 1–23.
- [4] V. G. KAC, P. CHEUNG. *Quantum Calculus*. Universitext, Springer-Verlag, New York, 2002.
- [5] CH. POMMERENKE. *On meromorphic starlike functions*. Pacific J. Math., 1963, **13**: 221–235.
- [6] J. E. MILLER. *Convex meromorphic mapping and related functions*. Proc. Amer. Math. Soc., 1970, **25**: 220–228.
- [7] H. M. SRIVASTAVA, K. SUCHITHRA, B. A. STEPHEN, et al. *Coefficient bounds for certain classes of meromorphic functions*. J. Inequal. Pure Appl. Math., 2008, 1–9.
- [8] R. M. ALI, V. RAVICHANDRAN. *Classes of meromorphic α -convex functions*. Taiwanese J. Math., 2010, **14**(4): 1479–1490.
- [9] M. K. AOUF. *Coefficient results for some classes of meromorphic functions*. J. Natur. Sci. Math., 1987, **27**(2): 81–97.
- [10] S. R. KULKARNI, S. S. JOSHI. *On a subclass of meromorphic univalent functions with positive coefficients*. J. Indian Acad. Math., 2002, **24**(1): 197–205.
- [11] V. KARUNAKARAN. *On a class of meromorphic starlike functions in the unit disc*. Math. Chronicle, 1976, **4**(2-3): 112–121.
- [12] Wancang MA, D. MINDA. *A Unified Treatment of Some Special Classes of Univalent Functions*. Int. Press, Cambridge, MA, 1994.
- [13] Z. NEHARRI. *Conformal Mapping*. McGraw-Hill, New York, 1952.
- [14] A. O. MOSTAFA, M. K. AOUF, H. M. ZAYED, et al. *Convolution conditions for subclasses of meromorphic functions of complex order associated with basic Bessel functions*. J. Egyptian Math. Soc., 2017, **25**(3): 286–290.
- [15] M. K. AOUF, A. O. MOSTAFA, H. M. ZAYED. *Convolution properties for some subclasses of meromorphic functions of complex order*. Abstr. Appl. Anal., 2015, Art. ID 973613, 6 pp.